

BUY AT STORE - TWO DAY BOWEL PREP - SAME DAY



The following instructions are your physician's specific instructions. Please follow the instructions carefully to ensure a successful prep.

You can reach Your Patient Advisor with non-medical prep questions at: **800.349.0285**.

You can reach your physician's office through United Digestive's Practice Network at **866.468.6242**.

Special Notes: If you are on aspirin, Coumadin (Warfarin), Plavix (clopidogrel), Xarelto (rivaroxaban) Eliquis (apixaban) or other blood thinning medication, follow the instructions provided by your gastroenterologist regarding if/when they should be discontinued before your procedure. If you are unsure, call your gastroenterologist.

PREP WEEK AT A GLANCE

<p>7 days to procedure</p>	<p>Review prep instructions thoroughly and the medication changes/restrictions starting today.</p> <p> STOP taking iron supplements and multivitamins with iron, high dose of fish oil, and phentermine.</p> <p>Purchase the following over the counter items at a pharmacy or grocery store:</p> <ul style="list-style-type: none">a. Two 28 oz bottles of Gatorade or Propel/G2/other sugar free electrolyte drink if you are diabetic (no red, purple, orange, or blue)b. One 8.3 oz bottle of MiraLAX powder (238 grams/14 dose)c. One box of Dulcolax tablets (5mg) <p>DIABETIC PATIENTS If you are a diabetic patient, please reference the additional diabetic instructions in this packet, located on page 5.</p>
<p>5 days to procedure</p>	<p> STOP eating any raw vegetables or vegetables containing seeds, corn, popcorn, nuts, and seeds. In addition, stop taking any fiber supplements until after the procedure.</p> <p>Confirm arrangements with your driver.</p>
<p>2 Days to procedure</p>	<p>Between 4:00 pm and 7:00 pm two (2) days before your procedure, you may have a light dinner (example: soup, sandwiches, eggs). No steak, potatoes, pasta, or salad. Otherwise, you may start your clear liquid diet.</p> <p>After your light meal, begin a clear liquid diet. NO SOLID FOOD, MILK OR DAIRY.</p> <p>Prep at a Glance</p> <ul style="list-style-type: none">● 7:00 pm, take 2 Dulcolax tablets. See step-by-step instructions located on p 2.
<p>1 day to procedure</p>	<p>NO SOLID FOODS, CLEAR LIQUIDS ONLY upon rising, until after your procedure. A clear liquid diet is necessary for a colonoscopy.</p> <p>Prep at a Glance</p> <ul style="list-style-type: none">● 10:00 am, take 2 Dulcolax tablets.● 4:00 pm, drink first dose of the prep solution.● 11:00 pm, drink second dose of the prep solution.● Drink plenty of water and liquids throughout the day to avoid dehydration.
<p>Procedure Day</p>	<ul style="list-style-type: none">● Nothing by mouth after 12:30 am (midnight), or the procedure may be rescheduled.● NO GUM OR HARD CANDY till after your procedure. You may brush your teeth but do not swallow.● If you take daily prescribed medications, please reference p 3 for specific instructions.

BUY AT STORE - TWO DAY BOWEL PREP - SAME DAY

Clear Liquid Diet Details: **NO RED** or **PURPLE, ORANGE, or BLUE**. **NO DAIRY**

Approved

Sodas, coffee, tea, clear juices, fitness waters, Popsicles without pulp, chicken, vegetable, & beef broth, gelatin

Avoid

No milk/dairy
No juices with pulp
NO RED or **PURPLE**
NO ORANGE or **BLUE**



Prep Day: Two (2) days before your procedure

Between 4:00 pm and 7:00 pm two (2) days before your procedure, you may have a light dinner (example: soup, sandwiches, eggs). No steak, potatoes, pasta, or salad. Otherwise, you may start your clear liquid diet.


After your light meal, begin a clear liquid diet. **NO SOLID FOOD, MILK OR DAIRY**. Drink plenty of water throughout the day to avoid dehydration.

Step

1

7:00 PM Two (2) days before your procedure:

Take 2 (5mg) Dulcolax tablets with a glass of water.



Prep Day: The day before your procedure

Clear liquids only the day prior to the procedure. No solid foods upon rising, until after your procedure. Drink plenty of water throughout the day to avoid dehydration.

- Discontinue alcohol one day prior to procedure.
- Diabetic Patients** – Take ½ of your usual diabetic medication (oral/injectable) dose today.

Step

2

10:00 AM the day prior to your procedure:

Take 2 (5mg) Dulcolax tablets with a glass of water.



Step

3

3:00 PM the day prior to your procedure:

Pour the 56 oz of Gatorade or other electrolyte drink into a pitcher and chill in the refrigerator. (2-Bottles of 28 oz each)



Step

4

3:00 PM the day prior to your procedure:

Mix the **ENTIRE BOTTLE** (8.3 oz/238g) of MiraLAX with **56 oz** of chilled Gatorade or other electrolyte drink and continue to chill in the refrigerator.



BUY AT STORE - TWO DAY BOWEL PREP - SAME DAY

Step 5

4:00 PM the evening prior to your procedure:

Consume an 8 oz glass of the chilled prep mixture every 15 – 30 minutes until you have finished drinking approximately **HALF OF THE PREP MIXTURE (32 oz)**.

This process may take 1 hour.



Step 6

5:00 PM – 11:00 PM the day prior to your procedure:

Drink an additional **FOUR (4)** 8 oz glasses of allowed clear liquids **before 11:00 PM**.



Individual responses to laxatives may vary. This preparation will cause multiple bowel movements, so it is advised to stay close to a bathroom.

Step 7

11:00 PM the evening prior to your procedure:

Consume the **OTHER HALF OF THE PREP MIXTURE (32 oz)** by drinking an 8 oz glass of the solution every 15 – 30 minutes **until the prep is finished**.

You must finish the entire prep, even if your bowels are already clear.



FINISH PREP BY 12:30 AM. NOTHING BY MOUTH UNTIL AFTER YOUR PROCEDURE.

The day of your procedure

Medication Instructions

- For diabetic patients, hold all diabetic medications, including insulin, till after your procedure.
- Unless your physician gives you other instructions, prescribed medication(s) for asthma, blood pressure, heart, seizure, or anti-psychotic medications may be taken as scheduled with a small sip of water at least 2 hours prior to your procedure. Please do not take anything else by mouth until your procedure is completed.

BUY AT STORE - TWO DAY BOWEL PREP - SAME DAY

Additional Information

- Please refrain from smoking on the day of your procedure to comply with anesthesia requirements. If you are a smoker, it is important that you avoid smoking before your procedure.
- Asthma patients: please bring your inhaler with you.
- **12:30 AM – THE NIGHT BEFORE YOUR PROCEDURE**
 - NOTHING BY MOUTH
 - NOTHING TO EAT OR DRINK UNTIL AFTER YOUR PROCEDURE
 - NO GUM OR HARD CANDY
 - YOU MAY BRUSH YOUR TEETH BUT DO NOT SWALLOW
- If your procedure requires sedation, a driver (age 18 or over) must accompany you to your appointment. The person who is driving you home must remain at the facility while your procedure is being performed so that they are available in the unlikely event of an emergency. You may not go home in a taxi, rideshare (Uber/Lyft), or take public transportation. The sedation you will receive will prevent you from driving safely or working the day of your procedure. Following your procedure, do not drive, operate heavy machinery, or sign legal documents for 24 hours.
- **If you have any questions or concerns about the preparation, please contact Your Patient Advisor by calling 800.349.0285 or emailing support@yourpatientadvisor.com.**

It is important to leave your name for your Patient Advisor and verify your call back phone number if prompted. Your Patient Advisor will return your call as soon as possible.

DIABETIC PATIENT INSTRUCTIONS

It is important to keep your blood sugar controlled when you are having an outpatient procedure. The following directions will guide you in managing your diabetes medicines before your procedure.

THE DAY BEFORE YOUR PROCEDURE

- Take ½ of your usual diabetic medication (oral/injectable) dose today.

THE EVENING BEFORE YOUR PROCEDURE

- If you take Insulin:
 - Take half of your usual amount of evening insulin (see examples below which include all types of insulin, such as NPH, 70/30, 50/50, Regular, Humalog, and Lente).
 - Example 1: Your usual evening dose is 15 units of NPH and 8 units of regular. You would take 7 units of NPH and 4 units of regular the evening before your procedure.
 - Example 2: Your usual evening dose is 27 units of 70/30 insulin. You would take 13 units of 70/30 insulin the evening before your procedure.
- NOTHING TO EAT OR DRINK AFTER MIDNIGHT. This includes hard candy and chewing gum. You are allowed to brush your teeth, but do not swallow.

THE MORNING OF YOUR PROCEDURE

- Nothing to eat or drink. This includes hard candy and chewing gum. You are allowed to brush your teeth, but do not swallow.
- Check your blood sugar first thing in the morning (and at 12:00 noon if you are not at the hospital or outpatient facility) and write down the results. Bring these results and your blood sugar logbook with you (along with your diabetes pills and/or insulin) to the hospital or outpatient facility.
- If you are on diabetes pills or take insulin injections:
DO NOT take your pills or insulin the morning of your procedure.
- If you use an insulin pump:
 - Discontinue the bolus function four (4) hours before your arrival time.
 - If your blood sugar is in an acceptable range, decrease the basal function to 50% of your normal rate.
 - If your blood sugar is less than 60, take one of the following 2 hours before your arrival at the hospital or outpatient facility:
 - a. 3 glucose tablets
 - b. 4 oz. juice (clear juice such as apple or white grape)
 - c. 6 Life Savers™ (NO RED, PURPLE, ORANGE OR BLUE)

Bowel Prep Frequently Asked Questions

What is a clear stool?

A clear stool can have a slight tint of yellow or brown. It will be completely transparent and will not contain any solid matter.

I am not having bowel movements, what should I do?

Bowel movements can take up to 5-6 hours after beginning the prep to start. Be patient, continue to drink clear liquids. If you have not had a bowel movement by midnight the night prior to your procedure, you will need to reach the on-call physician for further instructions.

The prep is making me nauseous, what should I do?

If you develop nausea or vomiting, slow down the rate at which you drink the solution. Please attempt to drink all of the laxative solution even if it takes you longer. If vomiting persists, or you are not able to finish the preparation, stop the preparation and contact your physician's office during normal business hours or speak to the on-call physician.

What are some high fiber foods I should avoid?

Raw fruits and vegetables are typically high in fiber, as well as nuts, seeds, whole grain breads and beans and lentils.

What are some good options for low fiber foods?

Choose white bread and white rice for lower fiber options, as well as pastas made with white flour. Chicken, fish, dairy and eggs are also low in fiber and good choices for foods 2-3 days before you begin your prep.

If I eat popcorn or seeds 3 days before my procedure, do I need to reschedule?

You will not need to reschedule your procedure; however, the seeds or nuts may cause a difficulty in screening and require a need for rescreening. If you have eaten a large amount of seeds or nuts, you may want to contact your provider or their medical assistant.

Can I drink ALCOHOL on the liquid diet?

Alcohol is not allowed as part of the liquid diet.

Can I continue to be on the liquid diet after I begin consuming the laxatives?

Yes, you may continue the liquid diet until you are directed to discontinue anything by mouth, which is typically 4-6 hours prior to the procedure.

Why do I have to wake up so early for the 2nd dose, can't I take it all the night before? *(Not applicable for same day prep instructions)*

A prep that is taken at two different time periods is called a split-dose prep. This prep protocol has proven to be the most effective for a successful colonoscopy. It is essential that you follow the directions provided with your prep medications.

If I weigh under 100 pounds, do I need to take all the prep?

The liquid amount is not weight dependent. It is important to finish the prep for a successful colonoscopy.