

SUFLAVE™ Instructions

Your doctor has prescribed Suflave Bowel Prep Kit for your upcoming procedure. You may be feeling a little anxious, but don't worry, this instruction sheet will guide you through every step.

Access video
instructions on
our website.



Medication Guidelines

GLP-1 and phentermine medications

- Seven (7) days prior to your procedure, discontinue oral and injectable GLP-1 agonist, phentermine, and any weight loss medications, such as Bydureon BCise® (exenatide), Byetta® (exenatide), Mounjaro® (tirzepatide), Ozempic® (semaglutide), Rybelsus® (semaglutide), Saxenda® (liraglutide), Soliqua® (insulin glargine/lixisenatide), Trulicity® (dulaglutide), Victoza® (liraglutide), Wegovy® (semaglutide), Zepbound (tirzepatide), Adipex, Fastin®, Phentercot, Pro-Fast, Qsymia®.
- Discontinue fiber and iron supplements, multivitamins with iron and high doses of fish oil.

Blood Thinning Medications

- Your GI provider will advise you whether to stop taking any blood thinning medications (anticoagulants and antiplatelets), after consulting with your prescribing physician.
- If you are currently taking aspirin, you may continue your dosage.

Diabetic Medications

- Three (3) days prior, diabetic patients, do not take SGLT2 medications, such as JARDIANCE® (empagliflozin), INVOKANA® (canagliflozin), FARXIGA® (dapagliflozin), SYNJARDY® (empagliflozin/metformin), or STEGLUJAN® (ertugliflozin/sitagliptin), etc.

Diabetic Patients Discontinuing GLP or SGLT2 Medications

- Please monitor your blood sugar levels while off your medication and consult your prescribing physician to determine if an alternative agent is required in the interim.

Clear Liquid Diet Guidelines

- For a successful bowel cleansing prep, it is essential that you **purchase** clear liquid items as indicated.
- Purchase your clear liquid diet items. For comfort, purchase diaper rash ointment, your preferred two-or-three-ply toilet paper, and/or flushable wet wipes.

Allowed Items

- ✓ Water
- ✓ Clear broths
- ✓ Coffee / tea without milk or creamers
- ✓ Clear juices without pulp
- ✓ Gatorade®
- ✓ Jell-O®
- ✓ Pulp-free Popsicles®

Items to Avoid

- ✗ Dairy
- ✗ Anything with pulp
- ✗ Alcohol
- ✗ Solid food
- ✗ Red, orange, or purple juices

Three Days Before Your Procedure

- Stop eating raw vegetables or vegetables containing seeds, corn, popcorn, nuts, and seeds.

One Day Before Your Procedure

- Unless otherwise directed, **begin your clear liquid diet** in the morning and refrain from consuming solid food until after your procedure.
- For patients with **constipation**, please take 15 mg of Dulcolax (Bisacodyl) by mouth at 6 PM, along with your bowel prep.



- Prepare your SUFLAVE the evening before your procedure at 5:00 p.m. Pour one SUFLAVE packet into the bottle provided and fill with lukewarm water to the fill line. Mix well by shaking the bottle until dissolved and refrigerate for one hour.
- At 6:00 p.m., start drinking 8 ounces of solution every 15 minutes until the bottle is empty.
- After you drink the SUFLAVE, drink at least one additional 16-ounce glass of water over the next hour. You will begin feeling the effects soon, so it is recommended to be near a restroom
- Medication instructions for the evening before
 - Insulin dependent patients may take insulin if necessary, but at **half** the usual dose with close monitoring of glucose. Keep a sugary clear liquid available if necessary.
 - Non-insulin dependent diabetic patients, take your oral diabetic medication dose as usual.

The Day of Your Procedure

- To comply with anesthesia requirements, do not use any tobacco products or marijuana on the day of your procedure. This includes smokeless tobacco, chewing tobacco, vapes, e-cigarettes, ZYN pouches, etc.

6 hours before the procedure



- Start the second dose of SUFLAVE 6 hours before the procedure start time, following the same instructions as your first dose.
- After your second dose, be sure to drink an additional 16-ounce glass of water. You need to finish the second SUFLAVE dose plus an additional glass of water.
- You must finish the entire prep even if your bowels are already clear.
- You must finish your prep 4 hours before your procedure.

Medication instructions for the day of procedure

- Diabetic patients, follow these medication instructions.
 - Insulin dependent patients may take insulin if necessary, but at **half** the usual dose with close monitoring of glucose. Keep a sugary clear liquid available **if necessary** to regulate your blood sugar. Nothing by mouth 2 hours prior to your procedure.
 - Non-insulin dependent diabetic patients, **hold** your oral diabetic medication until after your procedure.
- Unless your doctor gave you other instructions, prescribed medication(s) may be taken as scheduled with a small sip of water at least 2 hours before your procedure. Do not take anything else by mouth until your procedure is completed.
- Asthma patients, bring your inhaler to your appointment.
- It is imperative that you take all seizure and blood pressure medication on the morning of the procedure.

4 hours before the procedure

- Nothing by mouth until after your procedure.
- Do not chew gum or eat hard candy.
- You may brush your teeth but do not swallow.

Additional Day of Instructions

Because your procedure requires sedation, a driver (age 18 or over) must accompany you to your appointment. The person who is driving you home must remain at the facility while your procedure is being performed so that they are available in the unlikely event of an emergency. You may **not** go home in a taxi, rideshare (Uber/Lyft), or take public transportation. The sedation you will receive will prevent you from driving safely or working the day of your procedure. Following your procedure, do not drive, operate heavy machinery, or sign legal documents for 24 hours.



Follow any additional instructions from your doctor.
If you have questions, you can contact Gifthealth.

For additional information, scan the QR code for FAQs.

Need further assistance?

- Chat now on www.gifthealth.com
- 833.614.2051
- care@gifthealth.com